

Analysis of Thermo elastic Behavior of Semi Infinite Rectangular Beam Due to Heat Generation

A. A. Kulkarni; R.T. Walde; A. A. Navlekar and N. W. Khobragade

1, 2. Research Scholar, Department of Mathematics, RTM, Nagpur University, Nagpur- 440 033,

3. Assist. Professor, Department of Mathematics, Pratishthan Mahavidyalaya, Paithan, Dist.Aurangabad- 431107, India.

4. Professor, Department of Mathematics, RTM, Nagpur University, Nagpur- 440 033, India

Abstract-This paper is concerned, with the determination of temperature distribution and thermal deflection at any points of the plate with the stated boundary conditions. The transient heat conduction equation is solved by using the finite Fourier cosine and Marchi-Fasulo transform techniques. The results are obtained in the form of infinite series.

Index Terms – Transient thermoelastic problem, temperature distribution, thermal stresses, semi-infinite rectangular beam.

I. INTRODUCTION

Adams and Bert [1], Tanigawa and Komatsubara [4] and Vihak et.al [5] have studied the direct problem of the thermo elasticity in a rectangular plate under thermal shock. Khobragade et.al. [2,6,7] has studied the inverse steady state thermo elastic problem to determine the temperature, displacement function and thermal stresses at the boundary of a thin rectangular plate. They have used the finite Fourier sine transform techniques. Khobragade and Lamba [3] have studied three dimensional coupled thermo elastic response of infinitely long hollow circular cylinder due to axisymmetric heating, considered under the thermo-mechanical coupling effect. This approach is based upon expressions for both temperature and the stress distribution and are determined from field equation of motion. Numerical calculations are carried out and results are depicted graphically.

In the present paper, an attempt has been made to determine the temperature distribution, displacement function and thermal stresses of a thin rectangular beam occupying the region $D : -a \leq x \leq a ; -b \leq y \leq b, 0 \leq z \leq \infty$ with known boundary conditions. Here Marchi-Fasulo transforms and Fourier cosine transform techniques have been used to find the solution.

II. STATEMENT OF THE PROBLEM

Consider a thin rectangular plate occupying the space $D : -a \leq x \leq a ; -b \leq y \leq b, 0 \leq z \leq \infty$ The displacement components u_x and u_y u_z in the x and y and z directions respectively as Tanigawa et.al [4] are

$$u_x = \int_{-a}^a \left[\frac{1}{E} \left(\frac{\partial^2 U}{\partial y^2} + \frac{\partial^2 U}{\partial z^2} - \nu \frac{\partial^2 U}{\partial x^2} \right) + \lambda T \right] dx \quad (1)$$

$$u_y = \int_{-b}^b \left[\frac{1}{E} \left(\frac{\partial^2 U}{\partial z^2} + \frac{\partial^2 U}{\partial x^2} - \nu \frac{\partial^2 U}{\partial y^2} \right) + \lambda T \right] dy \quad (2)$$

$$u_z = \int_0^\infty \left[\frac{1}{E} \left(\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} - \nu \frac{\partial^2 U}{\partial z^2} \right) + \lambda T \right] dz \quad (3)$$

Where E, ν , and λ are the young's modulus, Poisson's ratio and the linear coefficient of the thermal expansion of the material of the beam respectively and U (x,y,z,t) is the Airy's stress functions which satisfy the differential equation:

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) U(x, y, z, t) = -\lambda E \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \times T(x, y, z, t) \quad (4)$$

where T(x,y,z,t) denotes the temperature of a rectangular beam satisfy the following differential equation:

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} + \frac{g(x, y, z, t)}{k} = \frac{1}{\alpha} \frac{\partial T}{\partial t} \quad (5)$$

Where K is the thermal conductivity and α is the thermal diffusivity of the material, subject to initial condition

$$T(x, y, z, 0) = f(x, y, z) \quad (6)$$

The boundary conditions are

$$\left[T(x, y, z, t) + k_1 \frac{\partial T(x, y, z, t)}{\partial x} \right]_{x=a} = f_1(y, z, t) \quad (7)$$

$$\left[T(x, y, z, t) + k_2 \frac{\partial T(x, y, z, t)}{\partial x} \right]_{x=-a} = f_2(y, z, t) \quad (8)$$

$$\left[T(x, y, z, t) + k_3 \frac{\partial T(x, y, z, t)}{\partial y} \right]_{y=b} = f_3(x, z, t)$$

$$\left[T(x, y, z, t) + k_4 \frac{\partial T(x, y, z, t)}{\partial y} \right]_{y=-b} = f_4(x, z, t) \quad (9)$$

$$(10)$$

$$\left. \frac{\partial T(x, y, z, t)}{\partial z} \right|_{z=0} = 0 \quad (11)$$

$$\left. \frac{\partial T(x, y, z, t)}{\partial z} \right|_{z=\infty} = 0 \quad (12)$$

The stress components in terms of U(x, y, z, t) are given by

$$\sigma_{xx} = \left[\frac{\partial^2 U}{\partial y^2} + \frac{\partial^2 U}{\partial z^2} \right] \quad (13)$$

$$\sigma_{yy} = \left[\frac{\partial^2 U}{\partial z^2} + \frac{\partial^2 U}{\partial x^2} \right] \quad (14)$$

$$\sigma_{zz} = \left[\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} \right] \quad (15)$$

The equations (1) to (15) constitute the mathematical formulation of the problem under consideration.

III. SOLUTION OF THE PROBLEM

Applying finite Marchi-Fasulo transform (twice) defined in [] and Fourier cosine transform to the equations, we have

$$\frac{d\bar{T}_c}{dt} + \alpha q^2 \bar{T}_c = \frac{\alpha g_c}{k} + \Psi$$

This is a linear equation whose solution is given by

$$\bar{T}_c(m, n, \eta, t) = \int_0^t \left[\frac{\alpha g_c}{k} + \Psi \right] e^{-\alpha q^2(t-t')} dt' \quad (16)$$

Where, $\phi = \frac{Q_n(a)}{k_1} f_1 - \frac{Q_n(a)}{k_2} f_2$

, $\Phi = \frac{P_m(b)}{k_3} f_3 - \frac{P_m(-b)}{k_4} f_4$, $\phi + \Phi = \Psi$

Now, applying inversion of Fourier Cosine transform and finite Marchi-Fasulo transform to the equation (16), one obtains the expression for temperature distribution as

$$T(x, y, z, t) = \frac{2\eta}{\pi} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m(y)}{\mu_m} \right) \Lambda(z) \quad (17)$$

Where,

$$\Lambda(z) = \int_0^{\infty} B(t) \cos(\eta z) dz$$

$$B(t) = \int_0^t \left[\frac{\alpha g_c}{k} + \Psi \right] e^{-\alpha q^2(t-t')} dt'$$

which is the required solution.

IV. AIRY'S STRESS FUNCTIONS

Substituting the value of temperature distribution T(x,y,z,t) from (16) in equation (4)

one obtains

$$U(x, y, z, t) = -\frac{2\eta\pi E}{\pi} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m(y)}{\mu_m} \right) \Lambda(z) \quad (18)$$

V. DISPLACEMENT COMPONENTS

Substituting the values of Airy's stress function from equation (18) in the equation (1) to (3), one obtains

$$u_x = -\frac{2\eta\lambda}{\pi} \Lambda(z) \int_{-a}^a \left[\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m''(y)}{\mu_m^2} \right) - \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \eta^2 \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m(y)}{\mu_m} \right) \right] dx \quad (19)$$

$$u_y = -\frac{2\eta\lambda}{\pi} \Lambda(z) \int_{-b}^b \left[\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \eta^2 \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m(y)}{\mu_m} \right) - \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n''(x)}{\lambda_n^2} \right) \left(\frac{P_m(y)}{\mu_m} \right) \right] dy + \nu \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m''(y)}{\mu_m^2} \right) + \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m(y)}{\mu_m} \right) \quad (20)$$

$$u_z = -\frac{2\eta\lambda}{\pi} \Lambda(z) \int_0^{\infty} \left[\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n''(x)}{\lambda_n^2} \right) \left(\frac{P_m(y)}{\mu_m} \right) + \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m''(y)}{\mu_m^2} \right) \right] dz + \nu \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \eta^2 \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m(y)}{\mu_m} \right) - \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m(y)}{\mu_m} \right) \quad (21)$$

VI. DETERMINATION OF STRESS FUNCTION

Substituting the value of Airy's stress function U(x,y,z,t) from equation (18) in the equation (13) to (15) one obtains the stress functions as,

$$\sigma_{xx} = -\frac{2\eta\lambda E}{\pi} \Lambda(z) \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left[\sum_{m=1}^{\infty} \left(\frac{P_m''(y)}{\mu_m^2} \right) - \eta^2 \sum_{m=1}^{\infty} \left(\frac{P_m(y)}{\mu_m} \right) \right] \quad (22)$$

$$\sigma_{yy} = -\frac{2\eta\lambda E}{\pi} \Lambda(z) \sum_{m=1}^{\infty} \left(\frac{P_m(y)}{\mu_m} \right) \left[\sum_{n=1}^{\infty} \left(\frac{Q_n''(x)}{\lambda_n^2} \right) - \eta^2 \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \right] \quad (23)$$

$$\sigma_{zz} = -\frac{2\eta\lambda E}{\pi} \Lambda(z) \left[\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n''(x)}{\lambda_n^2} \right) \left(\frac{P_m''(y)}{\mu_m} \right) + \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left(\frac{Q_n(x)}{\lambda_n} \right) \left(\frac{P_m''(y)}{\mu_m^2} \right) \right] \quad (24)$$

VII. MATERIAL PROPERTIES

The numerical calculations has been carried out for an Aluminum (pure) rectangular beam with the material properties as,

Density $\rho = 169 \text{ lb/ft}^3$

Specific heat = 0.208 Btu/lb OF

Thermal conductivity $K = 117 \text{ Btu/(hr. ft OF)}$

Thermal diffusivity $\alpha = 3.33 \text{ ft}^2/\text{hr.}$

Poisson ratio $\nu = 0.35$

Coefficient of linear thermal expansion $\alpha_t = 12.84 \times 10^{-6} / \text{F}$

Lame constant $\mu = 26.67$

Young's modulus of elasticity $E = 70 \text{ G Pa}$

VIII. DIMENSIONS

The constants associated with the numerical calculation are taken as

Length of rectangular beam $x = 3 \text{ ft}$

Breath of rectangular beam $y = 2 \text{ ft}$

Height of rectangular beam $z = 10^3 \text{ ft}$

IX. CONCLUSION

The temperature distribution displacements and thermal stresses at any point of a thin rectangular object have been obtained, when the boundary conditions are known with the aid of finite Marchi-Fasulo transform and semi-infinite Fourier cosine transform techniques. The results are obtain in the form of infinite series.

ACKNOWLEDGEMENT

The authors are thankful to University Grant Commission, New Delhi for providing the partial financial assistance under major research project scheme.

REFERENCES

[1] Adams, R.J and Bert, C. W: Thermoelastic vibrations of a laminated rectangular plate subjected to a thermal Shock, Journal of thermal stresses, vol.22 (1999), pp. 875-895.

[2] Durge M. H. and Khobragade N.W.: An inverse unsteady-state thermo elastic problem of a thick rectangular plate, Bull. Of the cal. Math. Soc., 95,(6) PP. 497-500 (2003)

[3] Khobragade, N. W and Lamba N.K. : Analysis of coupled thermal stresses in a axi symmetric Hollow cylinder, International Journal of latest trend in Mathematics, Vol.1, issue2, PP.29-32 (2012)

[4] Tanigawa Y and Komatsubara, Y: Thermal stress analysis of a rectangular plate and its Thermal stress intensity factor

for compressive stress field, journal of thermal stresses, Vol.20 (1997). PP.517

[5] Vihak, V. M; Yuzvyak, M. Y and Yasinskij, A. V: The solution of the plane Thermoelasticity problem for a rectangular domain, Journal of thermal stresses, vol.21 (1998), pp. 545-561.

[6] Khobragade, N.W.; Hiranwar Payal; Roy, H.S. and Khalsa Lalsingh: Thermal Deflection of a Thick Clamped Rectangular Plate, vol.3 (2013), pp.346-348.

[7] Roy, H.S.; Bagade, S.H. and Khobragade, N.W.: Thermal Stresses of a Semi Infinite Rectangular Beam. vol.3 (2013), pp.442-445.




Fig. 1. Temperature distribution vs radius




Fig. 2. Radial Stresses vs radius




Fig. 3. Axial Stresses vs radius

different capacities. At present he is working as Professor. Achieved excellent experiences in Research for 15 years in the area of Boundary value problems and its application. Published more than 180 research papers in reputed journals. Fourteen students awarded Ph.D Degree and four students submitted their thesis in University for award of Ph.D Degree under their guidance.

AUTHOR BIOGRAPHY



Mrs. A.A. Kulkarni For being M.Sc in maths, she has been teaching since 1990 for 21 years at PCE of Maths, RTM Nagpur University, Nagpur and successfully handled different capacities.



Mr. R.T. Walde For being M.Sc in Maths, he is a research scholar of Ph.D. He is teaching since 2004 for 09 years at Junior college level, Nagpur.



Dr. N.W. Khobragade For being M.Sc in statistics and Maths he attained Ph.D. He has been teaching since 1986 for 27 years at PGTD of Maths, RTM Nagpur University, Nagpur and successfully handled