

A New Algebraic Method for Computing the Pfaffian

González, H.E.¹, and Carmona, L., J.J.²

¹ Information Technology Department, ININ, ITTLA

Information Technology Department, ININ

Abstract— In this paper we postulate a new decomposition theorem of a matrix $A \in \mathbb{R}^n$ into two matrices, namely, a lower triangular one, M , all whose entries are determinants, and an upper triangular matrix U whose entries are also in determinant form. Applying this new linear transformation to skew-symmetric matrix $A \in \mathbb{R}^{2n}$ by means of partial pivoting strategy, we obtain the Pfaffian of A from the principal diagonal of matrix U , namely:

$$Pf(A) = \frac{\prod_{k=2}^n u_{k,k}}{\prod_{l=1}^{n-1} u_{l,l}}$$

Furthermore, if we apply this new linear transformation in Jordan version with the above pivot strategy to the augmented matrix $(A|I) \in \mathbb{R}^{2n}$, we obtain a “Pfaffian adjugate” G matrix. With it, we can solve systems of skew-symmetric linear equations in general, analogously to the “Cramer’s Rule”, which can be seen as a ratio of two Pfaffians. Both algorithms present an $O(n^3)$ computational complexity.

Index Terms— Pfaffian, LU decomposition, Pfaffian-Cramer rule, Pfaffian-Adjugate Matrix.

I. INTRODUCTION

The determinant of a skew-symmetric matrix ($A = -A^T$) is the square of another expression, which is called the Pfaffian [1]. For example,

$$\begin{vmatrix} 0 & a_{12} & a_{13} & a_{14} \\ -a_{12} & 0 & a_{23} & a_{24} \\ -a_{13} & -a_{23} & 0 & a_{34} \\ -a_{14} & -a_{24} & -a_{34} & 0 \end{vmatrix} = (a_{12}a_{34} - a_{13}a_{24} + a_{14}a_{23})^2$$

This property was proved by Cayley in 1849 [2]. The determinant of a skew-symmetric matrix of odd order is always 0.

Recently, Knuth [3], in a short history of Pfaffians, argues that these are in some way more fundamental than determinants, to which they are closely related. Johann Friedrich Pfaff introduced the functions that now bear his name in 1815 [1, pp. 396-401] while developing a general method to solve systems of first-order partial differential equations.

He gave two procedures for listing all perfect matching’s (subset of edges which covers all vertices of the graph), and observed that, when the matching are ordered lexicographically, the corresponding signs are strictly alternating $+, -, +, \dots, +$.

Formally speaking, the Pfaffian of a skew-symmetric matrix $A \in \mathbb{R}^{2n}$ can be defined as follows [4]:

$$Pf(A) = \sum_{\text{perfect matchings } M} \text{sign } M \cdot \text{weight}(M) \quad (1)$$

Here, a perfect matching M with $k = \frac{n}{2}$ edges is written as

$$M = (i_1, j_1), (i_2, j_2), \dots, (i_k, j_k) \quad (2)$$

Where, by convention, $i_k < j_k$ for each k . The sign of the matching M is defined as the sign of

$$\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & \dots & n-1 & n \\ i_1 & j_1 & i_2 & j_2 & \dots & i_k & j_k \end{pmatrix}$$

When this is regarded as a permutation of $\{1, 2, \dots, n-1, n\}$. The weight of M is $(i_1, j_1) \cdot (i_2, j_2) \cdot \dots \cdot (i_k, j_k)$.

Because of their close connection with matching’s, Pfaffians are of considerable interest in combinatorial analysis, as seen in [5,6]. For example, for certain graphs, including the planar ones, it is possible to count the number of perfect matching’s in a polynomial time by using Pfaffians.

The relation

$$|A| = Pf \begin{pmatrix} 0 & A \\ -A^T & 0 \end{pmatrix}$$

Shows that determinants are just special cases of Pfaffians that correspond to bipartite graphs. In this sense, the Pfaffian is a more basic notion, which would deserve a more thorough understanding, despite the traditional prevalence of the determinant in the curricula and in applications. This work expects to contribute in that direction.

A Linear System of Equations (LSE) can be defined as a set of m equations with n unknowns represented by a matrix A , a vector b and an unknown vector x , namely, $Ax = b$. The Gaussian Transformation (GT) for solving an LSE has proved to be the best option for most practical applications. The new transformation proposed here can be obtained from

Manuscript received: 23 September 2018
 Manuscript received in revised form: 24 October 2018
 Manuscript accepted: 04 November 2018
 Manuscript Available online: 10 November 2018

it. Next, we briefly review this topic.

II. LU- MATRICIAL DECOMPOSITION WITH GT. NOTATION AND DEFINITIONS

The problem of solving a linear system of equations $\mathbf{Ax}=\mathbf{b}$ is central to the field of matrix computation. There are several ways to perform the elimination process necessary for its matrix triangulation. We will focus on the Doolittle-Gauss elimination method: the algorithm of choice when \mathbf{A} is square, dense, and un-structured.

Let us assume that $\mathbf{A} \in \mathbf{R}^{n \times n}$ is nonsingular and that we wish to solve the linear system $\mathbf{Ax}=\mathbf{b}$. Here we show how for an exact arithmetic and partial pivoting and column interchanges some Gauss transformations $\mathbf{M}_1, \dots, \mathbf{M}_{n-1}$ can nearly always be found such that $\mathbf{M}_{n-1} \dots \mathbf{M}_2 \mathbf{M}_1 \mathbf{A} = \mathbf{U}$ is upper triangular [7]. The original $\mathbf{Ax}=\mathbf{b}$ problem is then equivalent to the upper triangular system $\mathbf{Ux}=(\mathbf{M}_{n-1} \dots \mathbf{M}_2 \mathbf{M}_1) \mathbf{b}$ which can be solved through back-substitution.

Suppose, then, that $\mathbf{A} \in \mathbf{R}^{n \times n}$ and that, for some $k < n$,

We have determined the Gauss transformations

$\mathbf{M}_1, \dots, \mathbf{M}_{k-1} \in \mathbf{R}^{n \times n}$ Such that

$$\mathbf{A}^{(k-1)} \equiv \mathbf{M}_{k-1} \dots \mathbf{M}_1 \mathbf{A} = \begin{pmatrix} \mathbf{A}_{11}^{(k-1)} & \mathbf{A}_{12}^{(k-1)} \\ \mathbf{0} & \mathbf{A}_{22}^{(k-1)} \end{pmatrix} \begin{matrix} (k-1) \\ (n-k+1) \end{matrix}$$

where: $\mathbf{A}_{11}^{(k-1)}$ is an upper triangular matrix.

$$\mathbf{A}_{22}^{(k-1)} = \begin{pmatrix} a_{kk}^{(k-1)} & \dots & a_{kn}^{(k-1)} \\ \vdots & \ddots & \vdots \\ a_{nk}^{(k-1)} & \dots & a_{nn}^{(k-1)} \end{pmatrix}$$

Now, if

and $a_{kk}^{(k-1)} \neq 0$, then the multipliers :

$$m_i = \frac{a_{ik}^{(k-1)}}{a_{kk}^{(k-1)}}; i = k+1, \dots, n$$

defined.

So, we have the following

Definition. An elementary lower triangular matrix of order n and index k is a matrix of the form [8]

$$\mathbf{M}_k = \mathbf{I}_n - \mathbf{m} \mathbf{e}_k^T$$

where:

$$\mathbf{e}_k^T = (0, \dots, 0, 1, 0, \dots, 0)^T$$

$$\mathbf{m}^T = (0, 0, 0, \dots, 0, m_{k+1}, \dots, m_n)$$

$\leftarrow k\text{-times}$

In general an elementary lower triangular matrix has the above form.

The computational significance of elementary lower triangular matrices is that they can be used to introduce zero components into a vector. Thus,

$$\mathbf{M}_k \cdot \begin{pmatrix} a_{11} \\ \vdots \\ a_{k1} \\ a_{k+1,1} \\ \vdots \\ a_{n1} \end{pmatrix} = \begin{pmatrix} a_{11} \\ \vdots \\ a_{k1} \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

The matrix \mathbf{M}_k is said to be a GT. The vector \mathbf{m} is referred to as the Gauss vector. The components of \mathbf{m} are known as multipliers.

Then it follows that

$$\mathbf{A}^{(k)} \equiv \mathbf{M}_k \mathbf{A}^{(k-1)} = \begin{pmatrix} \mathbf{A}_{11}^{(k)} & \mathbf{A}_{12}^{(k)} \\ \mathbf{0} & \mathbf{A}_{22}^{(k)} \end{pmatrix} \begin{matrix} (k) \\ (n-k) \end{matrix}$$

Where $\mathbf{A}_{11}^{(k)}$ is an upper triangular matrix.

This process illustrates the k -th step of the decomposition process, in which we used

$$(\mathbf{M}_k \dots \mathbf{M}_1)^{-1} = \mathbf{M}_1^{-1} \dots \mathbf{M}_k^{-1} = \prod_{i=1}^k (\mathbf{I}_n + \mathbf{m}^{(i)} \mathbf{e}_i^T) = \mathbf{I}_n + \sum_{i=1}^k \mathbf{m}^{(i)} \mathbf{e}_i^T$$

We find the final expression for the decomposition process as

$$\mathbf{A} = \begin{pmatrix} \mathbf{L}_{11}^{(k)} & \mathbf{0} \\ \mathbf{L}_{21}^{(k)} & \mathbf{I}_{n-k} \end{pmatrix} \begin{pmatrix} \mathbf{I}_k & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_{22}^{(k)} \end{pmatrix} \begin{pmatrix} \mathbf{A}_{11}^{(k)} & \mathbf{A}_{12}^{(k)} \\ \mathbf{0} & \mathbf{I}_{n-k} \end{pmatrix} \Rightarrow$$

$$(\mathbf{M}_k \dots \mathbf{M}_1)^{-1} \equiv \begin{pmatrix} \mathbf{L}_{11}^{(k)} & \mathbf{0} \\ \mathbf{L}_{21}^{(k)} & \mathbf{I}_{n-k} \end{pmatrix} = \mathbf{I}_n + (\mathbf{m}^{(1)}, \dots, \mathbf{m}^{(k)}, 0, \dots, 0).$$

In general, the forward elimination consists of $n-1$ steps. At the k -th step, multiples of the k -th equations are subtracted from the remaining equations to eliminate the k -th variable.

If the pivot element $a_{k,k}^{(k)}$ is null or “sufficiently small”, it is advisable to interchange equations before this is done through \mathbf{P} , a permutation matrix that records the row exchanges as detailed below.

$$U = MA = \begin{pmatrix} 1 & 0 \\ -a_{21} & a_{11} \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} \\ 0 & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{pmatrix}$$

$n=3$
 $k=1,2$

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}; M_1' = \begin{pmatrix} 1 & 0 & 0 \\ -a_{21} & a_{11} & 0 \\ -a_{31} & 0 & a_{11} \end{pmatrix};$$

$$U_1 = M_1' A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \\ 0 & \begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} \end{pmatrix};$$

$$M_2' = \frac{1}{a_{11}} \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{11} & 0 \\ 0 & -\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{pmatrix};$$

$$M = M_2' M_1' = \begin{pmatrix} 1 & 0 & 0 \\ -a_{21} & a_{11} & 0 \\ -\begin{vmatrix} a_{31} & a_{32} \\ a_{21} & a_{22} \end{vmatrix} & -\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{pmatrix};$$

The last row of the matrix **M** multiplied for the last column of the matrix **A** is equivalent to Laplace Expansion of **A** taking out the last column. Then we have $u_{33} = |A|$:

$$U = MA = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \\ 0 & 0 & \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \end{pmatrix};$$

n
 $k=1,2, \dots, n-1$

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1,n-1} & a_{1,n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2,n-1} & a_{2,n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3,n-1} & a_{3,n} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \dots & a_{n-1,n-1} & a_{n-1,n} \\ a_{n,1} & a_{n,2} & a_{n,3} & \dots & a_{n,n-1} & a_{n,n} \end{pmatrix};$$

$$M = M_{n-1}' \dots M_1' = \begin{pmatrix} m_{11} & 0 & 0 & \dots & 0 & 0 \\ m_{21} & m_{22} & 0 & \dots & 0 & 0 \\ m_{31} & m_{32} & m_{33} & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ m_{n-1,1} & m_{n-1,2} & m_{n-1,3} & \dots & m_{n-1,n-1} & 0 \\ m_{n,1} & m_{n,2} & m_{n,3} & \dots & m_{n,n-1} & m_{n,n} \end{pmatrix}$$

Where

$$m_{11} = 1; m_{21} = -a_{21}; m_{22} = a_{11};$$

$$m_{31} = -\begin{vmatrix} a_{31} & a_{32} \\ a_{21} & a_{22} \end{vmatrix}; m_{32} = -\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix}; m_{33} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix};$$

$$m_{n-1,1} = -\begin{vmatrix} a_{n,1} & a_{n,2} & a_{n,3} & \dots & a_{n,n-2} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2,n-2} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3,n-2} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n-2,1} & a_{n-2,2} & a_{n-2,3} & \dots & a_{n-2,n-2} \end{vmatrix};$$

$$m_{n-1,2} = -\begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1,n-2} \\ a_{n,1} & a_{n,2} & a_{n,3} & \dots & a_{n,n-2} \\ a_{3,1} & a_{3,2} & a_{3,3} & \dots & a_{3,n-2} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n-2,1} & a_{n-2,2} & a_{n-2,3} & \dots & a_{n-2,n-2} \end{vmatrix};$$

$$m_{n-1,3} = -\begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1,n-2} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2,n-2} \\ a_{n,1} & a_{n,2} & a_{n,3} & \dots & a_{n,n-2} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n-2,1} & a_{n-2,2} & a_{n-2,3} & \dots & a_{n-2,n-2} \end{vmatrix};$$

$$m_{n-1,n-1} = -\begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1,n-2} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2,n-2} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3,n-2} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n-2,1} & a_{n-2,2} & a_{n-2,3} & \dots & a_{n-2,n-2} \end{vmatrix}$$

$$m_{n,1} = \begin{vmatrix} a_{n,1} & a_{n,2} & a_{n,3} & \cdot & \cdot & \cdot & a_{n,n-1} \\ a_{21} & a_{22} & a_{23} & \cdot & \cdot & \cdot & a_{2,n-1} \\ a_{31} & a_{32} & a_{33} & \cdot & \cdot & \cdot & a_{3,n-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdot & \cdot & \cdot & a_{n-1,n-1} \end{vmatrix};$$

$$U = MA = \begin{pmatrix} u_{11} & u_{12} & u_{13} & \cdot & \cdot & \cdot & u_{1,n-1} & u_{1,n} \\ 0 & u_{22} & u_{23} & \cdot & \cdot & \cdot & u_{2,n-1} & u_{2,n} \\ 0 & 0 & u_{33} & \cdot & \cdot & \cdot & u_{3,n-1} & u_{3,n} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \cdot & \cdot & \cdot & u_{n-1,n-1} & u_{n-1,n} \\ 0 & 0 & 0 & \cdot & \cdot & \cdot & 0 & u_{n,n} \end{pmatrix}$$

Where

$$u_{11} = a_{11}; u_{12} = a_{12}; u_{13} = a_{13}; u_{1,n-1} = a_{1,n-1};$$

$$u_{1,n} = a_{1,n}$$

$$u_{22} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}; u_{23} = \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix};$$

$$u_{2,n-1} = \begin{vmatrix} a_{11} & a_{1,n-1} \\ a_{21} & a_{2,n-1} \end{vmatrix}; u_{2,n} = \begin{vmatrix} a_{11} & a_{1,n} \\ a_{21} & a_{2,n} \end{vmatrix}$$

$$u_{33} = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}; u_{3,n-1} = \begin{vmatrix} a_{11} & a_{12} & a_{1,n-1} \\ a_{21} & a_{22} & a_{2,n-1} \\ a_{31} & a_{32} & a_{3,n-1} \end{vmatrix};$$

$$u_{3,n} = \begin{vmatrix} a_{11} & a_{12} & a_{1,n} \\ a_{21} & a_{22} & a_{2,n} \\ a_{31} & a_{32} & a_{3,n} \end{vmatrix}$$

The Laplace Expansion of sub-matrix $A_{n-1,n-1}$ taking out the last column, is equivalent to multiply the $(n-1)$ -th row of the matrix M by the $(n-1)$ -th column of A . Then, we have

$$u_{n-1,n-1} = |A_{n-1,n-1}| \text{ and}$$

$$u_{n-1,n-1} = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \cdot & \cdot & \cdot & a_{1,n-1} \\ a_{21} & a_{22} & a_{23} & \cdot & \cdot & \cdot & a_{2,n-1} \\ a_{31} & a_{32} & a_{33} & \cdot & \cdot & \cdot & a_{3,n-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdot & \cdot & \cdot & a_{n-1,n-1} \end{vmatrix};$$

In a similar way, we have

$$u_{n-1,n} = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \cdot & \cdot & \cdot & a_{1,n} \\ a_{21} & a_{22} & a_{23} & \cdot & \cdot & \cdot & a_{2,n} \\ a_{31} & a_{32} & a_{33} & \cdot & \cdot & \cdot & a_{3,n} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdot & \cdot & \cdot & a_{n-1,n} \end{vmatrix};$$

Finally, having the last row of M multiplied by the last column of A , we have

$$m_{n,2} = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \cdot & \cdot & \cdot & a_{1,n-1} \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdot & \cdot & \cdot & a_{n,n-1} \\ a_{31} & a_{32} & a_{33} & \cdot & \cdot & \cdot & a_{3,n-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdot & \cdot & \cdot & a_{n-1,n-1} \end{vmatrix};$$

$$m_{n,3} = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \cdot & \cdot & \cdot & a_{1,n-1} \\ a_{21} & a_{22} & a_{23} & \cdot & \cdot & \cdot & a_{2,n-1} \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdot & \cdot & \cdot & a_{n,n-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdot & \cdot & \cdot & a_{n-1,n-1} \end{vmatrix};$$

$$m_{n,n-1} = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \cdot & \cdot & \cdot & a_{1,n-1} \\ a_{21} & a_{22} & a_{23} & \cdot & \cdot & \cdot & a_{2,n-1} \\ a_{31} & a_{32} & a_{33} & \cdot & \cdot & \cdot & a_{3,n-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdot & \cdot & \cdot & a_{n,n-1} \end{vmatrix};$$

$$m_{n,n} = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \cdot & \cdot & \cdot & a_{1,n-1} \\ a_{21} & a_{22} & a_{23} & \cdot & \cdot & \cdot & a_{2,n-1} \\ a_{31} & a_{32} & a_{33} & \cdot & \cdot & \cdot & a_{3,n-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdot & \cdot & \cdot & a_{n-1,n-1} \end{vmatrix}$$

and

$$u_{n,n} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & \dots & a_{1,n} \\ a_{21} & a_{22} & a_{23} & \dots & \dots & a_{2,n} \\ a_{31} & a_{32} & a_{33} & \dots & \dots & a_{3,n} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ a_{n,1} & a_{n,2} & a_{n,3} & \dots & \dots & a_{n,n} \end{pmatrix}$$

Now, if **M** is the new transformation, then we have

$$M = \prod_{k=n-1}^1 M'_k ; U = MA$$

In order to solve the linear system of equations **Ax = b**, we have

$$MAx = Mb$$

$$Ux = Mb$$

We can use the “backward process” and solve the linear system of equations using only determinants. For a matrix **A** with floating point entries this process requires

$$\frac{n(n-1)(2n-1)}{6} + \frac{(n-2)(n-1)}{2} = \frac{n^3}{3} - \frac{4}{3}n + 1$$

Floating point multiplications.

VI. NEW DECOMPOSITION THEOREM WITH PARTIAL PIVOTING

Gauss elimination on real numbers is generally unstable due to the possibility of finding arbitrarily small pivots. This process can be alleviated, however, by exchanging rows during the elimination. In our case, only when a pivot is zero, we exchange the row in a similar way to the Gauss process. The following theorem is given without proof:

Theorem. Let $A \in \mathbb{R}^{n \times n}$. Suppose that the New Transformation $M'_1 \dots M'_{k-1}$; row's permutation matrices $P = P_1 \dots P_{k-1}$ have been determined so that $U = M'_{k-1} P_{k-1} \dots M'_1 P A$. Then, the Upper Matrix **U** is obtained from **PA** without exchanging any rows and $U = MPA$. Furthermore, if *exch* \equiv number of row's exchanges, we have $|A| = (-1)^{exch} u_{n,n}$

VII. A NEW ALGEBRAIC METHOD FOR COMPUTING THE PFAFFIAN

Applying this new linear transformation to skew-symmetric matrix $A \in \mathbb{R}^{2n}$ by means of partial pivoting strategy, we obtain the Pfaffian of $A \in \mathbb{R}^{2n}$ from the principal diagonal of matrix **U**, namely:

$$Pf(A) = \frac{\prod_{k=2}^n u_{k,k}}{\prod_{l=1}^{n-1} u_{l,l}} \quad (5)$$

, where $(k=2,4,6,\dots,n)$ and $(l=1,3,5,\dots,n-1)$.

Proof. It follows from an induction on *n*:
n=2

$$A = \begin{pmatrix} 0 & a_{12} \\ -a_{12} & 0 \end{pmatrix}; P_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; P_1 A = \begin{pmatrix} -a_{12} & 0 \\ 0 & a_{12} \end{pmatrix}; M_1 = \begin{pmatrix} 1 & 0 \\ 0 & -a_{12} \end{pmatrix};$$

$$M_1 P_1 A = \begin{pmatrix} 1 & 0 \\ 0 & -a_{12} \end{pmatrix} \begin{pmatrix} -a_{12} & 0 \\ 0 & a_{12} \end{pmatrix} = \begin{pmatrix} -a_{12} & 0 \\ 0 & -a_{12}^2 \end{pmatrix};$$

If $M = M_1$ and $P = P_1$, then

$$U = MPA = \begin{pmatrix} 1 & 0 \\ 0 & -a_{12} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & a_{12} \\ -a_{12} & 0 \end{pmatrix} =$$

$$= \begin{pmatrix} 1 & 0 \\ 0 & -a_{12} \end{pmatrix} \begin{pmatrix} -a_{12} & 0 \\ 0 & a_{12} \end{pmatrix} = \begin{pmatrix} -a_{12} & 0 \\ 0 & -a_{12}^2 \end{pmatrix}$$

With this new method, the Pfaffian is the result of dividing the last value among the first of the main diagonal of the matrix **U**:

$$Pf(A) = \frac{-a_{12}^2}{-a_{12}} = a_{12}$$

n=4

$$A = \begin{pmatrix} 0 & a_{12} & a_{13} & a_{14} \\ -a_{12} & 0 & a_{23} & a_{24} \\ -a_{13} & -a_{23} & 0 & a_{34} \\ -a_{14} & -a_{24} & -a_{34} & 0 \end{pmatrix}; P_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix};$$

$$P_1 A = \begin{pmatrix} -a_{12} & 0 & a_{23} & a_{24} \\ 0 & a_{12} & a_{13} & a_{14} \\ -a_{13} & -a_{23} & 0 & a_{34} \\ -a_{14} & -a_{24} & -a_{34} & 0 \end{pmatrix}$$

$$M_1 P_1 A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -a_{12} & 0 & 0 \\ a_{13} & 0 & -a_{12} & 0 \\ a_{14} & 0 & 0 & -a_{12} \end{pmatrix} \begin{pmatrix} -a_{12} & 0 & a_{23} & a_{24} \\ 0 & a_{12} & a_{13} & a_{14} \\ -a_{13} & -a_{23} & 0 & a_{34} \\ -a_{14} & -a_{24} & -a_{34} & 0 \end{pmatrix} =$$

$$= \begin{pmatrix} -a_{12} & 0 & a_{23} & a_{24} \\ 0 & -a_{12}^2 & -a_{12}a_{13} & -a_{12}a_{14} \\ 0 & a_{12}a_{23} & a_{13}a_{23} & \begin{vmatrix} a_{13} & a_{34} \\ a_{12} & a_{24} \end{vmatrix} \\ 0 & a_{12}a_{24} & \begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix}^+ & a_{14}a_{24} \end{pmatrix};$$

Where $\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix}^+ = a_{14}a_{23} + a_{12}a_{34}$ is the *permanent*, similar to determinant, except that each term in the development is written with a plus sign.

Next:

$$M_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & a_{23} & a_{12} & 0 \\ 0 & a_{24} & 0 & a_{12} \end{pmatrix};$$

$$M_2 M_1 P_1 A = \begin{pmatrix} -a_{12} & 0 & a_{23} & a_{24} \\ 0 & -a_{12}^2 & -a_{12}a_{23} & -a_{12}a_{14} \\ 0 & 0 & 0 & a_{12} \left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right) \\ 0 & 0 & a_{12} \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right) & 0 \end{pmatrix};$$

$$P_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix};$$

$$P_2 M_2 M_1 P_1 A = \begin{pmatrix} -a_{12} & 0 & a_{23} & a_{24} \\ 0 & -a_{12}^2 & -a_{12}a_{23} & -a_{12}a_{14} \\ 0 & 0 & a_{12} \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right) & 0 \\ 0 & 0 & 0 & a_{12} \left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right) \end{pmatrix};$$

$$M_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & - \frac{\left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right)}{a_{12}} \end{pmatrix};$$

$$U = M_3 P_2 M_2 M_1 P_1 A = \begin{pmatrix} -a_{12} & 0 & a_{23} & a_{24} \\ 0 & -a_{12}^2 & -a_{12}a_{23} & -a_{12}a_{14} \\ 0 & 0 & a_{12} \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right) & 0 \\ 0 & 0 & 0 & \left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)^2 \end{pmatrix};$$

The Pfaffian is obtained starting from a process that could be described as a “backward simplification”. This is because it starts from the ultimate diagonal element of the matrix U , and then divided among the penultimate one and, in turn, among the antepenultimate one and, this way, until arriving to the first value of the diagonal:

$$Pf(A) = \frac{\left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)^2}{a_{12} \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right)} = \frac{\left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)^2}{-a_{12}^2 \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right)} = \frac{\left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)^2}{- \left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)} = a_{14}a_{23} + a_{12}a_{34} - a_{13}a_{24}$$

The Pfaffian can be expressed by *permanents* too:

$$Pf(A) = \frac{\left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)^2}{a_{12} \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right)} = \frac{\left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)^2}{-a_{12}^2 \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right)} = \frac{\left(- \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right) \right)^2}{\left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right)} = a_{14}a_{23} + a_{12}a_{34} - a_{13}a_{24}$$

On the other hand one has that:

$$\bar{M} = M_3 P_2 M_2 M_1 P_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -a_{12} & 0 & 0 & 0 \\ -a_{12}a_{24} & a_{12}a_{14} & 0 & -a_{12}^2 \\ a_{23} \left(\begin{vmatrix} a_{14} & a_{13} \\ a_{24} & a_{23} \end{vmatrix} + a_{12}a_{34} \right) & a_{13} \left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right) & a_{12} \left(\begin{vmatrix} a_{14} & a_{13} \\ a_{24} & a_{23} \end{vmatrix} + a_{12}a_{34} \right) & 0 \end{pmatrix};$$

$$U = \bar{M} A =$$

$$\begin{pmatrix} -a_{12} & 0 & a_{23} & a_{24} \\ 0 & -a_{12}^2 & -a_{12}a_{23} & -a_{12}a_{14} \\ 0 & 0 & a_{12} \left(\begin{vmatrix} a_{14} & a_{34} \\ a_{12} & a_{23} \end{vmatrix} - a_{13}a_{24} \right) & 0 \\ 0 & 0 & 0 & \left(\begin{vmatrix} a_{13} & a_{14} \\ a_{23} & a_{24} \end{vmatrix} - a_{12}a_{34} \right)^2 \end{pmatrix}$$

Finally, for the general case, we have:

$$A = \begin{pmatrix} 0 & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} & \dots & a_{1n} \\ -a_{12} & 0 & a_{23} & a_{24} & a_{25} & a_{26} & \dots & a_{2n} \\ -a_{13} & -a_{23} & 0 & a_{34} & a_{35} & a_{36} & \dots & a_{3n} \\ -a_{14} & -a_{24} & -a_{34} & 0 & a_{45} & a_{46} & \dots & a_{4n} \\ -a_{15} & -a_{25} & -a_{35} & -a_{45} & 0 & a_{56} & \dots & a_{5n} \\ -a_{16} & -a_{26} & -a_{36} & -a_{46} & -a_{56} & 0 & \dots & a_{6n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ -a_{1n} & -a_{2n} & -a_{3n} & -a_{4n} & -a_{5n} & -a_{6n} & \dots & 0 \end{pmatrix};$$

Then, for $(a_{00}^{(1)} = 1)$ and

$$k = 1, \dots, n-1$$

$$M_k =$$

$$\begin{pmatrix} 1 \\ \dots \\ \dots \\ 1_{k-1} \\ \dots \\ \dots \\ - \frac{a_{k+1}^{(k)}}{a_{k-1}^{(k)}} & \frac{a_k^{(k)}}{a_{k-1}^{(k)}} \\ \dots \\ \dots \\ - \frac{a_{nk}^{(k)}}{a_{k-1}^{(k)}} & \frac{a_{kk}^{(k)}}{a_{k-1}^{(k)}} \end{pmatrix}$$

$$\frac{\mathbf{A}^{\text{Adj}} \mathbf{Ax}}{\text{Pf}(\mathbf{A})} = \frac{\mathbf{A}^{\text{Adj}} \mathbf{b}}{\text{Pf}(\mathbf{A})}; \quad \frac{|\mathbf{A}| \mathbf{x}}{\text{Pf}(\mathbf{A})} = \mathbf{Gb}; \quad \mathbf{x} = \frac{\mathbf{Gb}}{\text{Pf}(\mathbf{A})} \quad (9)$$

This new result is a Pfaffian-Cramer-type solution in $O(n^3)$.

**EXAMPLE OF SOLUTION OF AX=B WITH THE G MATRIX :
VOLTERRA-LOTKA EQUATION FOR PREDATOR-PREY
DYNAMICS**

Suppose we have two competing species of fish, labeled prey (small fish) and predator (large fish). Let N_1 and N_2 denote the number of small and large fish, respectively. The differential equations satisfied by N_1 and N_2 , in the classic Volterra-Lotka model [12] are

$$\frac{dN_1}{dt} = \alpha_1 N_1 - \lambda_1 N_1 N_2 \quad (10)$$

$$\frac{dN_2}{dt} = -\alpha_2 N_2 + \lambda_2 N_1 N_2 \quad (11)$$

Where

α_1 = natural growth rate of small fish in the absence of large fish

λ_1 = death rate per encounter of small fish due to large fish

α_2 = natural death rate of large fishes in the absence of small fish

λ_2 = efficiency of turning predated small fishes due to predation

$-\lambda_1 N_1 N_2$ = gives the rate of small fish being lost

$\lambda_2 N_1 N_2$ = gives the growth rate of the population of large fish

For n species, the system of differential equations is in the form

$$\frac{dN_i}{dt} = k_i N_i + \frac{1}{\beta_i} \sum_{j=1}^n a_{ij} N_i N_j, \quad (i = 1, \dots, n) \quad (12)$$

Where

$a_{ij} = -a_{ji}$ and $a_{ii} = 0$ for all i ; the constants k_i and β_i are positive and are called equivalence numbers. Equations (12) are called the Volterra-Lotka equations, and play an important role in Ecology.

Suppose we are interested in the equilibrium solutions of system (12), in which case

$$\frac{dN_i}{dt} = 0 \quad \forall i \quad (13)$$

System (12) becomes

$$N_i \left(k_i \beta_i + \sum_{j=1}^n a_{ij} N_j \right) = 0$$

or, provided that $N_i \neq 0$ for $(i=1, \dots, n)$

$$\sum_{j=1}^n a_{ij} N_j = -k_i \beta_i, \quad (i = 1, \dots, n) \quad (14)$$

If the number of species is odd, $n = 2N+1$, no equilibrium solutions can exist, because a_{ij} is a skew-symmetric matrix of odd order.

Simple and interesting algebraic considerations permit us to study and describe the situation when the number of species is even, $n = 2N$.

Writing (14) in matrix notation, we have

$$\mathbf{Ax} = \mathbf{b} \quad (15)$$

Where \mathbf{A} is an even skew-symmetric matrix ($\mathbf{A} = -\mathbf{A}^T$) and \mathbf{x} and \mathbf{b} are $2N$ -dimensional vectors. Equation (15) is readily solved by the Pfaffian-Cramer's Rule, provided that $\text{Pf}(\mathbf{A}) \neq 0$

An order $2N$ Pfaffian is a triangular array of $N(2N-1)$ elements (i,j) :

$$Pf = \begin{vmatrix} a_{12} & a_{13} & a_{14} & \dots & a_{1,2N} \\ & a_{23} & a_{24} & \dots & a_{2,2N} \\ & & a_{34} & \dots & a_{3,2N} \\ & & & \dots & \dots \\ & & & & a_{2N-1,2N} \end{vmatrix} \quad (16)$$

which may be expanded by its first row in the same way as a determinant, except that the minor of an element (i,j) is a Pfaffian of order $2(N-1)$ obtained from Pf by deleting both the i -th row and column and the j -th row and column [13,14,15] Thus, if $n=4$ we have

$$Pf(\mathbf{A}) = \begin{vmatrix} a_{12} & a_{13} & a_{14} \\ & a_{23} & a_{24} \\ & & a_{34} \end{vmatrix} = a_{12} a_{34} - a_{13} a_{24} + a_{14} a_{23}$$

The Cramer's Rule reduces, in this case, to the ratio of two pfaffians. So, we will calculate the Pfaffians

$$Pf(\mathbf{A}_1) = \begin{vmatrix} -b_2 & -b_3 & -b_4 \\ & a_{23} & a_{24} \\ & & a_{34} \end{vmatrix} = -b_2 a_{34} + b_3 a_{24} - b_4 a_{23}$$

$$Pf(\mathbf{A}_2) = \begin{vmatrix} b_1 & a_{13} & a_{14} \\ & -b_3 & -b_4 \\ & & a_{34} \end{vmatrix} = b_1 a_{34} + b_4 a_{13} - b_3 a_{14}$$

$$Pf(\mathbf{A}_3) = \begin{vmatrix} a_{12} & b_1 & a_{14} \\ & b_2 & a_{24} \\ & & -b_4 \end{vmatrix} = -b_4 a_{12} - b_1 a_{24} + b_2 a_{14}$$

$$Pf(\mathbf{A}_4) = \begin{vmatrix} a_{12} & a_{13} & b_1 \\ & a_{23} & b_2 \\ & & b_3 \end{vmatrix} = b_3 a_{12} - b_2 a_{13} + b_1 a_{23}$$

Finally

$$\mathbf{x}_i = \frac{Pf(\mathbf{A}_i)}{Pf(\mathbf{A})} \quad (i = 1, \dots, 4) \quad (17)$$

Otherwise, we have

$$\mathbf{G} = \begin{pmatrix} 0 & -a_{34} & a_{24} & -a_{23} \\ a_{34} & 0 & -a_{14} & a_{13} \\ -a_{24} & a_{14} & 0 & -a_{12} \\ a_{23} & -a_{13} & a_{12} & 0 \end{pmatrix}$$

If $\mathbf{g}_i^T \in \mathbf{G}$ is the i -th row vector of \mathbf{G} , and $\mathbf{a}_i \in \mathbf{A}$ is the column vector of \mathbf{A} , we have

$$\mathbf{x}_i = \frac{\mathbf{g}_i^T \mathbf{b}}{\mathbf{g}_i^T \mathbf{a}_i} = \frac{\text{Pf}(\mathbf{A}_i)}{\text{Pf}(\mathbf{A})} \quad (i = 1, \dots, 4) \quad (18)$$

IX. CONCLUSION

In this paper we have introduced a new theorem on the decomposition into determinants of matrix $\mathbf{A} \in \mathbf{R}^n$ and the new linear transformations, expressed as equations (3) and (4). Most simultaneous linear equation systems can also be solved with these new linear transformations. The result is Cramer-type solutions in $O(n^3)$. Furthermore, we have obtained the Pfaffian of \mathbf{A} , expressed as equation (5) from the \mathbf{U} matrix.

On the other hand, we have proposed a modified Doolittle-Gauss-Jordan elimination process in two versions: the first one applied to the augmented matrix $(\mathbf{A}|\mathbf{I})$ and the second to the augmented matrix $(\mathbf{A}|\mathbf{b})$. The first version is an algorithm to calculate the new \mathbf{G} matrix from $\mathbf{A} \in \mathbf{R}^{2n}$, expressed as equation (8). The second version is a new direct method to solve linear system, equation (9), if $(\mathbf{A}, \mathbf{b}) \in \mathbf{R}^{2n}$. The above algorithms calculate Pfaffian-Cramer-type solutions of the linear systems.

Gaussian elimination is usually the most economical way to solve $\mathbf{Ax} = \mathbf{b}$. Nevertheless there is one reason why this new method might be relevant when $\mathbf{A} \in \mathbf{R}^{2n}$: while the Gaussian elimination is oriented to traditional prevalence of the determinants, this new method is focused to a pfaffian approach.

Finally, when referring to the Cramer's rule, it has been affirmed by G. Strang [11] that: "...Thus each component of x is a ratio of two determinants, a polynomial of degree n divided by another polynomial of degree n . This fact might have been recognized from Gauss elimination, but it never was". Not only has this fact been made evident in the present paper but also it has been shown that the Pfaffian-Cramer's Rule, is actually a ratio of two Pfaffians, derived from the Gauss elimination process.

REFERENCES

- [1] Muir, T. A treatise on the Theory of Determinants MacMillan and Co., London 1882 Repr. Dover, New York 1960.
- [2] Cayley, A. On the theory of permutants, Cambridge and Dublin Mathematical Journal 7 (1852), 40-51. Reprinted in his Collected Mathematical Papers 2, 16-26.
- [3] Knuth, D.E. Overlapping Pfaffians Electron. J. Comb. 3 (1996), No.2, article R5, 13, pp. Printed version: J. Comb. 3 (1996), No. 2, 147-159.

- [4] Rote, G. Division-Free Algorithms for the Determinant and the Pfaffian: Algebraic and Combinatorial Approaches Lectures of the Graduate Program Computational Discrete Mathematics, Berlin 1991-2000.
- [5] Lovász, L., Plummer, M.D. Matching Theory. Ann. Discr. Math., Vol. 29. North-Holland Mathematics Studies, Vol. 121. Amsterdam 1986.
- [6] Stembridge, J.R. Nonintersecting paths, pfaffians, and plane partitions. Adv. Math. 83 (1990), 96-113.
- [7] Householder, A.S. The theory of Matrices in Numerical Analysis Blaisdell Publishing Company, N.Y. First Edition, Second Printing Pages: 122-146 1965.
- [8] Stewart, G.W. Introduction to Matrix Computations Academic Press, Inc. Page: 115, 120 1973.
- [9] Golub, G.H., Van Loan, Ch. F. Matrix Computations John Hopkins University Press Page:56 1983.
- [10] Grosswald, E. Topics from the Theory of Numbers Mc Millan Company, N.Y. Pages.:277-279., 1966.
- [11] Strang, G. Linear Algebra and its Applications Academic Press, Inc. Pages: 163-164, 1976.
- [12] Caianiello, E.R. Combinatorics and Renormalization in Quantum Field Theory W.A. Benjamin, Inc. Reading, Massachusetts 1973.
- [13] Scott, R.F. and Mathews, G.B. The Theory of Determinants Cambridge University Press, N.Y. 2nd ed., pp. 92-97, 1904.
- [14] Muir, T. A treatise on the Theory of Determinants MacMillan and Company, Ltd., London pp. 196-203, 1882.
- [15] Caianiello, E.R. and Fubini, S. Nuovo cimento 9, 1218 (1952).



AUTHOR BIOGRAPHY

H. E. González. He received the PhD in Operations Research at the National Autonomous University of Mexico (UNAM) in 2005. He has been working in research activities for more than twenty years. He has published a book and more than ten scientific papers. Presently, he is a full-time researcher at the National Institute of Nuclear Research (ININ) at the Department of Systems.



J.J. Carmona L., He received his B.S. in Electronic Engineering from UAM. Presently, he is a full-time researcher at the National Institute of Nuclear Research (ININ) at the Department of Systems.